

STIC-ILL

From: Baskar, Padmavathi
Sent: Monday, June 23, 2003 12:33 PM
To: STIC-ILL
Subject: 09/300612, please consider this as rush

*Miss
only*

Bone "The Pathogenesis of Sepsis" Ann Int Med 115(6) 457-469 1991.

Guyton "Textbook of Medical Physiology" 8th Ed 269-271 1991.

Mousstach e et al "Mechanisms of Resistance of the Opossum to Some Snake Venoms" Toxicon 17(Suppl. 1) 130 1979

Domont et al. "Natural Anti-Snake Venom Proteins" Toxicon 29(10) 1183-1194 1991.

Perales et al "Neutralization of the Oedematogenic activity of Bothrops Jararaca venom on the Mouse Paw by an antithroptic

Fraction Isolated from Opossum serum" Agents Actions 37(3-4) 250-259 1992.

Tomihara et al. "Purification of Three Antihemorrhagic Factors From The Serum of A Mongoose" Toxicon 25(6) 685-689 1987.

Perates et al. "Anti Snake Venom Protein from Didelphidae" Abstract 10th World Congress. Toxicon 30(5-6) 543 1992.

Menchaca et al. "The Purification & Characterization of An Antihemorrhagic Factor in Opossum Serum" Toxicon 19(5) 623-632 1981.

Toxicon 14(4) 337-340, 1976

Tarng et al , Toxicon 24(6) 567-573, 1986.

Toxicon 34(11-12) 1313-6, 1996.

Toxicon 36(10) 1451-9, 1998.

HIGH TECH SEPARATIONS NEWS 1996, V9,N4, SEP1996

Toxicon 37(6) 949-954, 1999.

Toxicon 37(5) 703-728, 1999.

biochimica et biophysica acta 1995, 1245 (2) 232-8

Padma Baskar
Art Unit 1645
Patent Examiner/Biotechnology
CM-1, 8E-13
703-308-8886

MECHANISMS OF RESISTANCE OF THE OPOSSUM TO SOME SNAKE VENOMS

H. Moussatché, A. Yates, F. Leonardi and L. Borche

Universidad Centro Occidental, Unidad de Investigación en Ciencias Fisiológicas and Sección de Fisiopatología, Escuela de Medicina de la U.C.O., Barquisimeto, Venezuela

Opossum (Didelphis, g.) of several species showed a natural resistance against the pharmacological action of some snake venoms. VELLARD (1949) observed the resistance of Peruvian species of Didelphis against the venom of snakes of the Crotalinae family. More recently WERNER and VICK (1977) showed the resistance of Didelphis virginiana against the envenomation by the venoms of Crotalus and Agkistrodon species.

Three mechanisms, at least, may be playing a role in the resistance of these animals to envenomation: 1) Presence in the serum of a compound neutralising the action of the venoms; 2) Low sensitivity of the animal receptors to the autopharmacological substances released by the venoms; 3) No release or release of low quantities of the autopharmacological agents responsible of the animal envenomation.

In the present abstract the presence of a serum component neutralising some of the pharmacological actions of the venoms of Crotalidae family is related:

1. The serum of Didelphis injected by intraperitoneal route (around 6 mg/protein = 0.1 ml serum) protects mice against the action of the venoms of Bothrops jararaca and Crotalus adamanteus (LD = 100 %).

2. The protective effect of the serum against the venom lasts up to 24 hours.

3. The serum has protective action against the necrosis induced in the foot of the rabbit by the injection of the venom of B. jararaca.

4. In mice the serum did not afford protection against the venom of Crotalus durissus terrificus. Nevertheless, the Opossum itself is resistant to this venom.

5. A fraction was isolated from the Didelphis serum, by chromatography and electrophoresis, which possessed the protective properties of the serum described above.

VELLARD, J. (1949) Public. Museum Histor. Natural. Univ. Nac. San Marcos, Lima, Año 1, No. 2.
WERNER, R.N. and VICK, J.A. (1977) Toxicon 15, 29.

Supported by a grant of the Research Council of the Centro Occidental University.